

Early measurements using W/Z in ATLAS

D. Prieur

STFC Rutherford Appleton Laboratory, Harwell Science & Innovation Campus, Didcot, OX11 0QX, UK

On behalf of the ATLAS Collaboration

The LHC experiments are close to collecting collision data. An overview of first physics measurements of the W and Z production cross-sections is presented. The electron and muon decay channels are considered. Emphasis will be given to data-driven approaches.

1. INTRODUCTION

The study of the W and Z production is a fundamental area for the ATLAS [1] early running. These processes are very well understood theoretically and with high expected counting rates they will provide stringent tests of QCD. They will help to understand the detector performance and be used for: calibration of the electromagnetic calorimeter (EM), alignment of the muon spectrometer system, as well as for extracting the lepton identification efficiencies. The first measurements will consist in determining the W/Z and W/Z +jets cross-sections [2]. Increased statistics will give access to fundamental electroweak parameters. In this note, the emphasis is put on the early cross-section measurements assuming a LHC peak luminosity of $\mathcal{L} = 10^{31} \text{ cm}^{-2}\text{s}^{-1}$.

2. W/Z CROSS-SECTION MEASUREMENT IN ELECTRON FINAL STATES

The reconstruction of electrons is based on clusters in the EM calorimeter, with a matching track in the Inner Detector [1]. The identification of isolated high- p_T electrons is then based on the shapes of the EM showers, and on track reconstruction information. Three sets of identification criteria with different degrees of tightness (Loose, Medium, Tight) are used [2]. The selection of $W \rightarrow e\nu$ events proceeds as follows. The trigger selects events with at least one electron candidate with $E_T > 20 \text{ GeV}$. The analysis procedure selects events with exactly one electron candidate satisfying $E_T > 25 \text{ GeV}$, $|\eta| < 1.37$ or $1.52 < |\eta| < 2.4$ and the Medium-type electron identification criteria. Then, the reconstructed missing transverse energy should satisfy $\cancel{E}_T > 25 \text{ GeV}$, and the transverse mass of the (l, ν) system should satisfy $M_T > 40 \text{ GeV}$. The resulting transverse mass distribution is shown in Fig. 1. Jet events constitute the largest background component. The jet production cross-section and fragmentation properties at the LHC are not well known and induce a significant uncertainty on the magnitude of this background. A data-driven method to monitor the jet background is applied and is presented in section 4. A MC study of this channel for $\int \mathcal{L} dt = 50 \text{ pb}^{-1}$ gave a measured cross-section of $\sigma = 20530 \pm 40(\text{stat}) \pm 1060(\text{syst}) \pm 2050(\text{lumi}) \text{ pb}$, to be compared with an input value $\sigma = 20510 \text{ pb}$ [3, 4].

The analysis of $Z \rightarrow ee$ selects at least two electron candidates with $E_T > 10 \text{ GeV}$ at the trigger level. The presence of two loosely identified isolated electrons with $E_T > 15 \text{ GeV}$ and $|\eta| < 2.4$ is then required. The resulting di-electron invariant mass distribution is shown in Fig. 1. As in the $W \rightarrow e\nu$ analysis, the jet background is estimated by a data-driven method. The signal and background fractions are estimated simultaneously, via a fit to both contributions. The signal is described by the convolution of a Breit-Wigner and a Gaussian resolution function, and the background, completely dominated by jet events, by an exponential function. A MC study of this channel for $\int \mathcal{L} dt = 50 \text{ pb}^{-1}$ gave a measured cross-section of $\sigma = 2016 \pm 16(\text{stat}) \pm 72(\text{syst}) \pm 202(\text{lumi}) \text{ pb}$, to be compared with an input value $\sigma = 2015 \text{ pb}$.




Figure 1: Left: Transverse mass distribution in the $W \rightarrow e\nu$ channel for signal and background, for $\int \mathcal{L}dt = 50 \text{ pb}^{-1}$. Right: Di-electron invariant mass distribution in the $Z \rightarrow ee$ channel, for signal and background, for $\int \mathcal{L}dt = 50 \text{ pb}^{-1}$.




Figure 2: Left: Transverse mass distribution in the $W \rightarrow \mu\nu$ channel, for signal and background, for $\int \mathcal{L}dt = 50 \text{ pb}^{-1}$. Right: Di-muon invariant mass distribution in the $Z \rightarrow \mu\mu$ channel, for signal and background, for $\int \mathcal{L}dt = 50 \text{ pb}^{-1}$.

3. W/Z CROSS-SECTION MEASUREMENT IN MUON FINAL STATES

The analysis of $W \rightarrow \mu\nu$ selects events with at least one muon candidate with $E_T > 20 \text{ GeV}$ at the trigger level. The events are further selected requiring exactly one muon track candidate, identified in the muon and inner detector tracking system, satisfying $|\eta| < 2.5$ and $p_T > 25 \text{ GeV}$. The energy deposited in the calorimeter, in a cone of radius $\Delta R = 0.4$ along the muon track, must not exceed 5 GeV . In addition the event has to satisfy $\cancel{E}_T > 25 \text{ GeV}$ and $M_T > 40 \text{ GeV}$. Figure 2 shows the corresponding W transverse mass distribution before the M_T cut. In contrast to the electron channels, the jet background is less important here. The dominant backgrounds come from $W \rightarrow \tau\nu$ and $Z \rightarrow \mu\mu$ events. These processes are well understood theoretically and can be safely estimated based on simulation. A MC study of this channel for $\int \mathcal{L}dt = 50 \text{ pb}^{-1}$ gave a measured cross-section of $\sigma = 20530 \pm 40(\text{stat}) \pm 630(\text{syst}) \pm 2050(\text{lumi}) \text{ pb}$, to be compared with an input value $\sigma = 20510 \text{ pb}$.

The $Z \rightarrow \mu\mu$ analysis uses the 10 GeV single muon trigger. The data sample is further reduced by requiring two offline tracks with opposite charges, in the muon spectrometer only, with $|\eta| < 2.5$ and $p_T > 20 \text{ GeV}$. The invariant mass of the muon pair is required to fulfil $|91.2 \text{ GeV} - M_{\mu\mu}| < 20 \text{ GeV}$. The corresponding invariant mass distribution before the mass cut is shown in Fig. 2. In this channel, the dominant background originates from $t\bar{t}$ events. The jet background is expected to be smaller, but is theoretically not well known. Other backgrounds are smaller, theoretically well known, and contribute negligibly to the overall background uncertainty. A MC study of this channel for $\int \mathcal{L}dt = 50 \text{ pb}^{-1}$ gave a measured cross-section of $\sigma = 2016 \pm 16(\text{stat}) \pm 64(\text{syst}) \pm 202(\text{lumi}) \text{ pb}$, to be compared with an input value $\sigma = 2015 \text{ pb}$.

4. DATA-DRIVEN BACKGROUND ESTIMATION FOR $W \rightarrow e\nu$

The principle is to measure the normalisation and shape of the jet background before the $\cancel{E}_T$ cut, in a sufficiently pure jet sample. This sub-sample is then used to evaluate the rejection of the $\cancel{E}_T$ cut, allowing a realistic estimation




Figure 3: Comparison of the jet background (points with error bars) and the fitted background (rectangles), for an integrated luminosity of $\int \mathcal{L} dt = 50 \text{ pb}^{-1}$.

of the jet background in the $W \rightarrow e\nu$ selection. The jet background control sample is selected using a single photon trigger with $E_T > 20 \text{ GeV}$, and subsequent calorimeter only based electron identification. Simulation studies show that these selections provide a sample almost entirely composed of jet events, even at high values of $\cancel{E}_T$, and that the shape of the $\cancel{E}_T$ distribution is identical, within the statistical precision, to that of the jet background in the $W \rightarrow e\nu$ sample (see Fig. 3). Above $\cancel{E}_T > 10 \text{ GeV}$, the slope can be described with the convolution of an exponential and a second degree polynomial function. After the subtraction of the estimated background from the signal sample, the analysis then proceeds by applying the $\cancel{E}_T$ selection mentioned above.

5. $W/Z + \text{JETS}$

The production of W/Z +jets events is an interesting measurement in itself. In addition this process is a background to many other Standard Model and beyond Standard Model physics channels. Furthermore these channels will test jet reconstruction techniques. Compared to the W/Z inclusive production, more statistics is needed and the analysis done is based on an integrated luminosity of 1 fb^{-1} . Selections are similar to the inclusive W/Z production, except that one, two or three jets with $E_T > 40 \text{ GeV}$ are required. At larger jet multiplicities, the dominant background arises from top quark events. The jet energy scale is the largest source of systematic error on the cross section. The initial uncertainty on the jet energy scale is expected to be 5-10%. A jet energy scale with precision better than 10 % is required to distinguish between the LO/NLO predictions of the different Monte-Carlo generators.

6. CONCLUSIONS

The prospects for the measurement of the W and Z boson cross-sections in ATLAS have been presented. In the four channels ($W \rightarrow e\nu$, $Z \rightarrow ee$, $W \rightarrow \mu\nu$, $Z \rightarrow \mu\mu$) considered, high purity samples have been achieved after standard selections (high p_T lepton identification, isolation and $\cancel{E}_T$ in the W final states) [2]. The jet background has been controlled and subtracted using data driven methods.

References

- [1] The ATLAS Collaboration, G. Aad et al., “The ATLAS Experiment at the CERN Large Hadron Collider”, 2008 JINST 3 S08003
- [2] The ATLAS Collaboration, “Expected Performance of the ATLAS Experiment, Detector, Trigger and Physics”, CERN-OPEN-2008-020, Geneva, 2008, to appear
- [3] T. Sjostrand, S. Mrenna and P. Skands, JHEP 05, 026 (2006)
- [4] K. Melnikov and F. Petriello, Phys. Rev. D74, 114017 (2006)